

Alain Ramanantsoanirina <sup>1</sup>, Julie Dusserre <sup>2</sup>, Suchit Shrestha <sup>3</sup>, and Folkard Asch <sup>3</sup>

<sup>1</sup>FOFIFA, Madagascar; <sup>2</sup>CIRAD, France; <sup>3</sup>University of Hohenheim, Germany

## Introduction


High altitude rice cultivation is constrained by a short vegetation period due to low temperatures and thus by the time the crop needs to complete its cycle. Climate change is assumed to result in a rise in mean temperatures of 2–5 degrees depending on the simulation scenario. Thus, rice cropping in higher altitudes may become more favorable as long as precipitation is not a limiting factor. In order to match rainy season with crop duration in higher altitudes rice genotypes are needed that possess an early vigor, a short duration and a certain degree of drought resistance.




## Conclusions

- ✓ High altitude conditions increase crop duration.
- ✓ Highest yield were obtained with the recommended sowing date in mid altitudes.
- ✓ Sterility is not influenced by minimum air temperature at booting.
- ✓ Cold tolerant varieties show less sterility under high altitude conditions.


## Results and Discussion



- ❖ Crop duration increases with altitude.
- ❖ The effect is less pronounced in mid altitudes.




- ❖ All varieties yielded best in mid altitudes when sown at the recommended date.



- ❖ Sterility is independent of temperature in low altitude.
- ❖ Sterility is lower in mid altitude conditions.

## Materials and Methods

- 10 varieties of upland rice were planted at 5 monthly staggering sowing dates in three locations.
- Plot size was 1 X 1 m with 20 X 20 cm spacing between plants.
- Daily values for minimum and maximum temperature were recorded.
- Different physiological and phenological stages were observed during crop cycle.
- Biomass, yield, and yield components were determined at maturity.



**Acknowledgements** This study is part of a large project addressing „Adaptation of African agriculture to climate change“ funded by the German Ministry for collaboration and development through GTZ/BEAF. More information can be found at [www.risocas.de](http://www.risocas.de). Collaborators in this study are the University of Hohenheim; Germany, WARDA; Benin, and CIRAD; France. FOFIFA, Madagascar as a co-partner institute heartily thanks funding organization and Collaborating institutions.